

# Publicly Funded Immunization Schedules for Ontario – June 2022

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

Routine Schedule: Children Starting Immunization in Infancy													
Vaccine	Age	2 Months	4 Months	6 Months	1 Year $\Phi$	15 Months	18 Months	4 Years	Grade 7	14 Years	24 Years	$\geq 34$ Years $\Psi$	65 Years
<b>DTap-IPV-Hib</b> Diphtheria, Tetanus, Pertussis, Polio, <i>Haemophilus influenzae</i> type b		◆	◆	◆			◆						
<b>Pneu-C-13</b> Pneumococcal Conjugate 13		◆	◆		◆								
<b>Rot-1</b> Rotavirus		▲	▲										
<b>Men-C-C</b> Meningococcal Conjugate C					◆								
<b>MMR</b> Measles, Mumps, Rubella					■								
<b>Var</b> Varicella						■							
<b>MMRV</b> Measles, Mumps, Rubella, Varicella								■					
<b>Tdap-IPV</b> Tetanus, diphtheria, pertussis, Polio								◆					
<b>HB</b> Hepatitis B									●				
<b>Men-C-ACYW</b> Meningococcal Conjugate ACYW-135									●				
<b>HPV-9</b> Human Papillomavirus									●				
<b>Tdap</b> Tetanus, diphtheria, pertussis										◆	◆		
<b>Td (booster)</b> Tetanus, diphtheria												◆ Every 10 years	
<b>HZ</b> Herpes Zoster													■
<b>Pneu-P-23</b> Pneumococcal Polysaccharide 23													■ / ◆
<b>Tdap</b> Tetanus, diphtheria, pertussis													◆ One dose in every pregnancy, ideally between 27-32 weeks of gestation
<b>Inf</b> Influenza													◆ Every year in the fall *

◆ - A single vaccine dose given by intramuscular injection  
 ■ - A single vaccine dose given by subcutaneous injection  
 ▲ - A single vaccine dose given by mouth  
 ● - Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-9 is a 2 dose series (see Table 10). Each vaccine dose is given by intramuscular injection  
 $\Phi$  - Given no earlier than the 1<sup>st</sup> birthday, and prior to 16 months of age

$\Psi$  - Once a dose of Tdap is given in adulthood (24 years of age), adults should receive Td boosters every 10 years thereafter  
 ■ - HZ is a 2 dose series (see Table 12) given by intramuscular injection  
 \* - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given  $\geq 4$  weeks apart. Children who have previously received  $\geq 1$  dose of influenza vaccine should receive 1 dose per season thereafter  
**Note:** A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)